

1126-14-102

Alexander Varchenko* (anv@email.unc.edu), Department of Mathematics, Chapel Hill, NC.

Elliptic dynamical quantum group $E_{\tau,h}(gl_2)$ and elliptic equivariant cohomology of the cotangent bundles of Grassmannians. Preliminary report.

The torus T equivariant elliptic cohomology defines a functor $Ell_T : \{T - \text{spaces } X\} \rightarrow \{\text{schemes}\}$. For example, for the cotangent bundle of a Grassmannian the scheme $Ell_T(T^*Gr(k, n))$ is some explicitly given sub-scheme of $S^k E \times S^{n-k} E \times E^n \times E^2$ with coordinates $t_1, \dots, t_k, s_1, \dots, s_{n-k}, z_1, \dots, z_n, h, \lambda$, where t_i, s_j correspond to the Chern roots of the two standard vector bundles over the Grassmannian, z, y correspond to the torus parameters, λ is the dynamical parameter also called the Kähler parameter, and E is an elliptic curve.

I will define a class of line bundles on the scheme $\cup_{k=0}^n Ell_T(T^*Gr(k, n))$ such that the operator algebra of the elliptic dynamical quantum group $E_{\tau,h}(gl_2)$ will act on sections of those line bundles (a generator of the operator algebra will send a section of such a line bundle to a section of possibly another line bundle). That construction is an analog of the Yangian $Y(gl_2)$ action on the direct sum $\oplus_{k=0}^n H_T^*(T^*Gr(k, n))$ of equivariant cohomology.

This is a joint work with G.Felder and R.Rimanyi. (Received January 07, 2017)