

1126-35-388

**Andrii Anikushyn** and **Michael Pokojovy\*** (michael.pokojovy@kit.edu). *Multidimensional Thermoelasticity for Nonsimple Materials – Well-Posedness and Long-Time Behavior.*

We consider a PDE system

$$\begin{aligned}\rho\ddot{u}_i &= (A_{iJKj}u_{j,K} - \beta_{Ji}\dot{\tau} - (C_{iJKLIj}u_{j,IL} + M_{iJKL\tau,L}),K)_{,J} - E(|\dot{u}|)\dot{u}_i, \\ a\ddot{\tau} &= -\beta_{Ki}\dot{u}_{i,K} + m_{IJ}q_{I,J} + M_{jLKl}u_{j,LKl} + K_{IJ}\tau_{,IJ}, \\ \kappa\dot{q}_i &= \dot{\tau}_{,i} - q_i\end{aligned}$$

in  $\Omega \times (0, \infty)$  subject to the boundary conditions

$$u_i = 0, \quad u_{i,J} = 0, \quad \tau = 0 \text{ in } \partial\Omega \times (0, \infty)$$

and the initial conditions

$$\begin{aligned}u_i(\cdot, 0) &= u_i^0, \quad \dot{u}_i(\cdot, 0) = \dot{u}_i^0, \\ \tau(\cdot, 0) &= \tau^0, \quad \dot{\tau}(\cdot, 0) = \dot{\tau}^0, \quad q_i(\cdot, 0) = q_i^0 \text{ in } \Omega.\end{aligned}$$

Under appropriate assumptions, well-posedness and uniform stability are obtained. In the absence of a frictional damping, the lack of asymptotic stability is shown. (Received January 17, 2017)